

ALKASPIRIN *Tablets*

Buffered aspirin.

Composition:

Each tablet contains:

- Acetyl salicylic acid 300 mg.
- Magnesium carbonate 100 mg.
- Aluminum glycinate 100 mg /tab.

ALkASPIRIN is a very well tolerated type of buffered aspirin " it exerts the analgesic, antipyretic. Anti-inflammatory , ant rheumatic. and ant platelet effects of Aspirin but without causing the stomach upset oftenly produced by the plain types. This latter advantage of ALKASPIRIN is provided by Magnesium carbonate and Aluminium glycinate the two non-systemic antacids which protect the gastric mucosa against Aspirin-induced irritation".

Indications and Dosage:

Notice:

Supervision.

As analgesic for mild to moderate pain in cases of headache, toothache , neuralgia, myalgia, mild injuries, postoperative pain (after the 2nd or 3rd day), dysmenorrhoea, postparturn pain. etc , and as analgesic-antipyretic in colds, influenza, and other febrile conditions.

- **Adults: 1 - 2 tablets to be repeated after 4- 6 hours if required.**
- **Children 2 years and over (see precautions below) : 1/3 tablet per year of age up to a maximum of 2 tablets at the age of 10 years and over. This dose can be repeated every 6 hours of necessary for not more than 48 hours.**

— *For arthritis (Osseo, Rheumatoid, etc)*

- Adults 2 tablets every 4 - - 6 hours according to the condition.
Juvenile rheumatoid arthritis :1 /3 tablet/kg. wt/day in divided doses every 4 - - 6 hours.

— *For rheumatic fever*

(Treatment should be continued for 4 - - 8 weeks depending on clinical course and ESR value)

- Adults : initially 4 tablets every 4 - 6 hours according to the condition, to be reduced after one week to 3 tablets every 4 - 6 hours.
- Children initially 1/3 tablet/kg. wt/day in divided doses every 4-6 hours to be reduced after two days to 1/4 tablet/kg. wt/day for 4 weeks (to be tapered after that).

- *For long term prophylaxis in thrombotic diseases*

- Cerebral Transient Ischaemic Attacks (T.I.A) in men : 2 tablets every 12 hours (or 1 tablets every 6 hours).
- Unstable angina and post-myocardial infarction : 1 tablet daily.

Packing:

Boxes of 24 tablets.