

DOLOMINE *Sterile injection (I.M.&I.V.)*

Ketorolac Tromethamine

Composition:

Each ampoule (1ml) contains:

- Ketorolac Tromethamine 15 mg.

Properties:

Ketorolac is a NSAID . ketorolac inhibits cyclo-oxygenase enzyme preventing prostaglandin formation, thus prevents the inflammation and produces rapid analgesic activity.

Usage & Indications:

. DOLOMINE, is indicated for the short term (max.5 days).

Ketorolac is indicated in :

- Painful inflammatory conditions, (short term treatment).
- Management of post-operative pain .

Side Effects:

- Rashes, pain at the site of injection, headache, drowsiness, dizziness, nausea, dyspepsia.
- gastrointestinal pain, diarrhea, constipation, flatulence, vomiting, Oedema, hypertension & stomatitis, sweating & purpura.

Dosage & Administration :



DOLOMINE THERAPY IS NOT TO EXCEED 5 DAYS.

Regimen To Single Administration:

IM Dosing :

- Patients < 65 years of age : one dose of 60 mg.
- Patients > 65 years of age , renal impaired and / or less than 50 kg of body weight : one dose of 30 mg.

IV Dosing :

- Patients < 65 years of age : one dose of 30 mg.
- Patients > 65 years of age , renal impaired and / or less than 50 kg body weight : one dose of 15 mg.

The recommended initial dose is one ampoule IM , followed by one to two ampoules every 4 - 6 hours. The IV injection should be given over at least 15 second. If combined therapy daily doses in adult should not exceed 90 mg, & in elderly and under weight patients the daily doses should not exceed 60 mg . In case patients with mild impaired renal function the daily doses should not exceed 60 mg.

Packing:

- A box of 3 ampoules (1ml).