

FLAGIMIDE *Tablets*

For treatment of amoebiasis

Composition:

Each tablets contains :

- Metronidazole 250 mg.
- Diloxanide furoate 250 mg.



Properties:

FLAGIMIDE contains the two active principles Metronidazole and Diloxanide furoate which act together to ensure a comprehensive and efficient treatment for amoebiasis. Metronidazole is a very effective agent against vegetative forms of *Entamoeba histolytica* (both intestinal and extra intestinal forms) and is also effective against the cyst forms.

On the other hand, Diloxanide furoate acts on intestinal amoebicide and has a potent action specially on the cyst forms of the fore - mentioned parasite, and so it adds to and complements the action of Metronidazole.

Indications:

- Acute amoebic dysentery.
- Chronic amoebiasis.
- Amoebic hepatitis and liver abscess.

Side Effects:

Occasionally occur in some patients and may

take the form of : metallic taste, gastrointestinal upsets, hypersensitivity reactions, headache, sedation or excitation, urinary disturbances, joint pains, flattening of T wave in the ECG mild Neutrogena or leukopenia.

Dosage:

Tablets are to be given with or after meals.

- Adults : 2 tablets 3 times daily for 5-10 days.

- Children 7-13 years : 1 tablets 3 times daily for 5-10 days .

packing:

Boxes of 20 tablets (2 strips x 10 tablets)