

# **FLATIDYL** *tablets & drops*

**Antiflatulent**


## **Composition:**

- Flatidyl chewable tablets, each tablet contains:

- Activated dimethicone [ simethicone ] 40 mg.
- Aerosil 2 mg.

- Flatidyl drops : each 1 ml contains:

- Simethicone emulsion 30 % 133.3 mg, eq . to 40 mg simethicone pure.



## **Properties:**

Simethicone is non - toxic silicone - based surface active agent with an efficient de-foaming action which relieves flatulence by dispersing and preventing the formation of the mucous surrounded by gas pockets in G. I. tract. It acts in the stomach and intestines to change the surface tension of the gas bubbles enabling it to coalesce, thus gas is freed and eliminated more easily .

## **Indications & Usage:**

- To relieve the painful symptoms of the excess gas in the digestive tract.
- As adjunctive therapy in the treatment of many conditions in which gas retention may be problem such as post operative gaseous distention - air swallowing - functional dyspepsia - peptic ulcer - spastic or irritable

colon or diverticulitis .

- To prepare for endoscopic , radiological or sonographic investigations of the stomach and or bowl.

### **Dosage & Administration:**

**Flatidyl chewable tablets:** (Tablets are thoroughly chewed and followed by some water).

- Average dose: 1 -2 tablets after meals and at bedtime .

- As diagnostic aid: 2 tablets to be given 2 - 3 days before the day of examination .

**Flatidyl drops:**

- 40 mg [ 30 - 35 drops ] four times daily after meal and at bedtimes .

### **Packing:**

- Flatidyl chewable tablets: Boxes of 30 tablets (3 strips x 10 tablets).

- Flatidyl drops: Plastic bottle [ with dropper ] contains 10 ml of the suspension.