

IBUGESIC

IBUPROFEN

Anti-inflammatory – analgesic & antipyretic

Composition:

- **IBUGESIC TABLETS** : sugar coated tablets containing 200,400 & 600 mg ibuprofen.
- **IBUGESIC SYRUP** :A flavored syrup contains 100 mg ibuprofen/ 5 ml.
- **IBUGESIC CREAM** :Available in 10 % ibuprofen cream.
- **IBUGESIC SUPPOSITORIES** : each suppository contains 500 mg ibuprofen.


properties:

IBUGESIC (ibuprofen) is non-steroidal anti-inflammatory agent used in painful inflammatory conditions. It is absorbed from the GIT & peak plasma concentration occurs about 1-2 hours after ingestion. Ibuprofen is extensively bound to plasma proteins & has a half-life of about 2 hours. It is rapidly excreted in the urine mainly as metabolites & their conjugates. About 1 % is excreted in urine as unchanged ibuprofen & about 14 % as conjugated ibuprofen.

Indications:

" **IBUGESIC** has Anti-inflammatory – analgesic & antipyretic properties "

IBUGESIC is used in the management of



mild to moderate pain in condition such as:

- **Rheumatoid arthritis, osteoarthritis, non-articular rheumatism & soft tissues injuries.**
- **Dysmenorrhoea, migraine, post-operative pain, Ankylosing spondylitis.**
- **Sprain & strain.**
- **As antipyretic, effective in reducing fever.**

Side Effects:

IBUGESIC is well tolerated & the main side effects are gastro-intestinal disturbance such as:

- **Abdominal discomfort, nausea, vomiting, and abdominal pain.**
- **In rare conditions, bleeding or activation of peptic ulcer may occur.**
- **Skin rash of various types may occur.**
- **Very rarely, toxic amblyopia may occur and on cessation of treatment, the patient recovered.**

Dosage & Administration:

Note:

before food.

1 - Dosage For Tablets:

IBUGESIC 200:2 tablets 3 times daily.

IBUGESIC 400: 1 tablet 3 times daily.

IBUGESIC 600: 1 tablet 2 times daily.

2 - Dosage for syrup:

20mg/kg - 40mg/kg. body weight in divided doses.

3 - Dosage for suppositories:

(2-3) suppositories daily.

4 - Dosage for Cream:

IBUGESIC Cream: To be applied 2-3 times daily with gentle massaging to enhance absorption.

Presentation:

- IBUGESIC 200 tablets: 3 strips x 10 tablets in a carton box.

- IBUGESIC 400 tablets: 3 strips x 10 tablets in a carton box.

- IBUGESIC 600 tablets: 3 strips x 10 tablets in a carton box.

- IBUGESIC Syrup: A glass bottle x 100ml in a carton box.

- IBUGESIC Suppositories: 6 supp. of 500 mg. Ibuprofen.

- IBUGESIC Cream: A compressible tube of 20 gm.