

Spasmopyralgin-M Tablets, Drops , Ampoules 2&5 ml & Suppositories

Antispasmodic, Analgesic and Antipyretic.

Composition:

1 - Tablets:

Each tablet contains: Camylofin 25 mg -
Analgin 255 mg.

2 - Drops:

Each 1 ml contains: Camylofin 25 mg -
Analgin 255 mg.

3 - Ampoules 2 & 5 ml:

Each 1 ml contains: Camylofin 10mg -
Analgin 258 mg.

4 - Suppositories:

For adult - For children: Camylofin 20 mg -
Analgin 250 mg.

Properties:

Spasmopyralgin-m (different forms) provides a powerful antispasmodic, analgesic and antipyretic effects, produced by its contents of the efficient antispasmodic "Camylofin " and the potent analgesic - antipyretic " Analgin " . Spasmopyralgin removes spasm and pain effectively.

Indications & Usage:

- Biliary, renal and intestinal colics .
- Dysmenorrhoea and cervical spasm.
- As antipyretic.
- Pains due to pancreatitis.

Contra-Indications:

- Hypersensitivity to any of the components.



- Blood diseases, rapid cardiac arrhythmia.
- Narrow angle glaucoma.
- Prostatic hypertrophy and obstructive bowel diseases.

Use in Pregnancy and Nursing Mothers:

Contra-indicated

Dosage & Administration:

1 - Tablets:

Adults: 1-2 tablet 1-3 times daily.

Children: (10 - 12 years) 1 tablet 1-3 times daily.

2 - Injectable forms :

Adults:

In severe attacks: 1 injection of 5 ml I.V. to be repeated if necessary.

In less severe attacks: 2-5 ml I.M.

Children:

doses are to be given as directed by the physician. To be given from 1-3 times daily.

3 - Drops:

Children: 6 months to 3 years: 5- 10 drops.

3 years to 6 years: 12-15 drops.

6 years to 12 years : 17-20 drops.

4 - Suppositories:

Adults: 1 adult suppository 2-3 times daily.

Children: (Use children suppositories)

Infants from 6 months : ½ suppository 2-3 times daily.

Children up to 5 years : 1 suppository 1-3 times daily.

Over 5 years : 1 suppository 2-3 times daily.

Packing:

- Tablets: Strips x 10 tablets.

- Drops: Bottles of 15 ml with dropper.
- Ampoules: Boxes of 5 ampoules of 2 ml or 5 ml.
- Suppositories: Boxes of 5 suppositories for adults or for children.

Storage:

- Tablets , Drops , Ampoules: Store at room temperature, away from light & humidity.
 - Suppositories: Store in a cool place.
- Keep out of the reach of children.